




Inclusion Criteria
Patients with idiopathic scoliosis

Exclusion
Patients with non-idiopathic scoliosis, see Complex Spine Algorithm



Idiopathic Scoliosis Spinal Fusion Overview

Background & Problem:

The Spine CCGs (Idiopathic Scoliosis Spinal Fusion and Complex Spine Algorithms) have not been updated in 4-5 years and there is new evidence that should be included. There is a lack of data available for Complex Spine patients due to variable and unreliable documentation practices.

Inclusion:

Patients with idiopathic scoliosis.

Exclusion:

Patients with non-idiopathic scoliosis, see *Complex Spine Algorithm*.

Outcome Measure:

- ☐ Unanticipated ICU transfers
- ☐ Length of stay (days)
- ☐ Allogenic transfusion rate and volume
- ☐ SSI rate
- ☐ Readmission rate

Process Measures:

- ☐ Hours of epidural
- ☐ Hours of PCA

Balancing Measures:

- ☐ Unplanned return to the OR



Contributors

Executive Sponsor:

Erik King

Clinical Leads:

John Grayhack
Jill Larson
Brett Lullo
Nicholas Burjek

Key Members:

Joseph Janicki
Eric Fuller
Brigid Marietta
Katherine Kennelly
Emily Beneturski
Elizabeth Klein
Diane Sheehan
Kerryn Oreilly
Mehul Raval
Matthew Rowland
Hubert Benzon
Alexander Froyshteter
David Krodel
Natalie Weatherred
Laura Fountain
Sarah Duggan
Daniela Doyle
Alison Thompson
Katie Tillhof

Key Members Continued:

Thomas Moran
Yvonne Johnson
Jamie Burgess
Craig Smith
Julie Borrowman
John Sarmiento
Kawana Mitchell
Wayne Z. Wellington
Mary lafelice
Mai Quad
Doreen Stein
Sameer Patel
Caitlin Li
Amanda Bonebrake
Joe Baker
Jennifer Lavin
Eileen Romano
Lindsey Vitosky
Michael Olsen
Carolyn Howell
Manisha Patel
Anna Wood
Kate Testa

Center for Quality and Safety:

Rachel Sotolongo
Naomi Sullivan
Abbey Studer

Version History: October 2018, April 2024

References

- Changoor, S. Giakas, A., Sacks, K., Asma, A., Land, R.S., Yorgova, P., Rogers, K., Gabos, P.G., & Shah, S.A. (2023). The Role of Liposomal Bupivacaine in Multimodal Pain Management following Posterior Spinal Fusion for Adolescent Idiopathic Scoliosis Faster and Farther with Less Opioids. *Spine*, 42 (2), 11-16. <https://doi.org/10.1097/BRS.0000000000004702>
- Gausden, E.B., Brusalis C.M., Audsi, R.A., Swarup, I., Fu, M., Dodwell, E., & Fabricant, P.D. (2020). Efficacy of antifibrinolytics in pediatric orthopedic surgery: a systematic review and meta-analysis. *Journal of Pediatric Orthopaedics B*, 29 (1) 97-104. <https://doi.org/10.1097/BPB.0000000000000599>
- Johnson, D.J., Johnson, C.C., Goobie, S.M., Nami, N., Wetzler, J.A., Sponseller P.D., & Frank, S.M. (2017). High-dose Versus Low-dose Tranexamic Acid to Reduce Transfusion Requirements in Pediatric Scoliosis Surgery. *Journal of Pediatric Orthopaedics*, 37 (8) 552-557. <https://doi.org/10.1097/BPO.0000000000000820>
- Halanski, M.A., Cassidy, J.A., Hetzel, S., Reischmann D., & Hassan, N. (2014). The Efficacy of Amicar Versus Tranexamic Acid in Pediatric Spine Deformity Surgery: A Prospective, Randomized, Double-Blinded Pilot Study. *Spine Deformity*, 2, 191-197. <https://doi.org/10.1016/j.jspd.2014.02.001>
- McNicol, E.D., Tzortzopoulou, A., Schumann, R., Carr, D.B., & Kalra, A. (2016). Antifibrinolytic agents for reducing blood loss in scoliosis surgery in children (review). *Cochrane Database of Systematic Reviews* (9) 1-63. <https://doi.org/10.1002/14651858.CD006883.pub3>
- Oetgen, M.E., Weinstein, S.L., Andras, L., Shah, S., & Sucato, D.J. (2020). Scoliosis Round Table: What are your Optimal Surgical Strategies for a Double Major Curve In Adolescent Idiopathic Scoliosis. *Journal of the Pediatric Orthopaedic Society of North America*, 2 (1).
- Shapiro, F., Zurakowski, D., & Sethna, N.F. (2007). Tranexamic Acid Diminishes Intraoperative Blood Loss and Transfusion in Spinal Fusions for Duchenne Muscular Dystrophy Scoliosis. *Spine*, 32 (20) 2278-2283. <https://doi.org/10.1097/BRS.0b013e31814cf139>